



TITLE:

ON THE GLOBAL EXISTENCE OF SOLUTIONS  
FOR THE DISCRETE BOLTZMANN EQUATION  
WITH LINEAR AND QUADRATIC  
TERMS(Generalized Functions and  
Differential Equations)

AUTHOR(S):

YAMAZAKI, MITSURU

---

CITATION:

YAMAZAKI, MITSURU. ON THE GLOBAL EXISTENCE OF SOLUTIONS FOR THE DISCRETE BOLTZMANN EQUATION WITH LINEAR AND QUADRATIC TERMS(Generalized Functions and Differential Equations). 数理解析研究所講究録 1996, 935: 143-150

ISSUE DATE:

1996-01

URL:

<http://hdl.handle.net/2433/60013>

RIGHT:

# ON THE GLOBAL EXISTENCE OF SOLUTIONS FOR THE DISCRETE BOLTZMANN EQUATION WITH LINEAR AND QUADRATIC TERMS

MITSURU YAMAZAKI

Dept. Mathematical Science, Univ. Tokyo

In this paper, we study the discrete Boltzmann equation in one-dimensional space with linear and quadratic terms. This system, which is more general than the usual one by the intervention of linear terms, describes the gas motion of molecules which take only a finite number ( $\#I < \infty$ ) of velocities  $c_i (i \in I)$  under the interactions between particles represented by the quadratic terms and also under the reflection of molecules at the inner wall of an infinite thin tube, represented by the linear terms which we treated in the papers [9], [10], [11], [12], [13], [14].

$$(B) \quad \begin{cases} \frac{\partial u_i}{\partial t} + c_i \frac{\partial u_i}{\partial x} = Q_i(u) + L_i(u), \\ u_i|_{t=0} = u_i^0(x), \quad i \in I, t \in \mathbb{R}_+, x \in \mathbb{R}^1 \end{cases}$$

where

$$\begin{aligned} Q_i(u) &= \sum_{j,k,\ell \in I} (A_{ij}^{k\ell} u_k u_\ell - A_{k\ell}^{ij} u_i u_j), \\ L_i(u) &= \sum_{k \in I} (\alpha_i^k u_k - \alpha_k^i u_i). \end{aligned}$$

The physical theory imposes to this system the natural conditions :

Conditions .—

$$(*) \quad \begin{cases} A_{ij}^{k\ell} \geq 0, \quad A_{ij}^{k\ell} = A_{ji}^{k\ell} = A_{ij}^{\ell k}, \\ A_{k\ell}^{ij} \neq 0 \Rightarrow i \neq j, \\ \alpha_i^k \geq 0 \quad \text{and} \quad \alpha_i^i = 0 \quad \text{for all} \quad i, k \in I. \end{cases}$$

This linear terms are more general than the ones which are obtained by considering solutions around constant stationary solutions  $(M_i)$  with  $Q_i(M) = 0$ , called constant Maxwellian. We suppose furthermore the condition of distinct velocities and the microreversibility condition for the quadratic terms.

**Condition vd.**—

$$i \neq j \rightarrow c_i \neq c_j$$

**Condition  $\mu rQ$ .**—

$$\sum_{k, \ell \in I} A_{k\ell}^{ij} = \sum_{k, \ell \in I} A_{ij}^{k\ell} \text{ for } \forall i, j \in I$$

We introduce the space  $\mathcal{B}(\mathbf{R} \times [0, T])$  defined as follows : for  $T < \infty$  fixed,

$$\begin{aligned} \mathcal{B}(\mathbf{R} \times [0, T]) &= \{u(x, t) : u_i^\sharp(x, t) \in C(0, T; L_{loc}^1(\mathbf{R})) \\ \text{such that } \sum_{i \in I} \int_K \sup_{t \in [0, T]} |u_i^\sharp(x, t)| dx &< \infty, \text{ for } \forall K \subset \subset \mathbf{R}\}, \end{aligned}$$

$$\text{where } u_i^\sharp(x, t) = u_i(x + c_i t, t).$$

**Definition.**— Suppose the condition (vd). Let be  $u_i^0 \in L_{loc}^1(\mathbf{R})$  and  $u \in \mathcal{B}(\mathbf{R} \times [0, T])$ . We say that  $u$  is a solution of the system (B) with Cauchy data  $(u_i^0)_i$  if and only if

$$u_i^\sharp(x, t) = u_i^0(x) + \int_0^t \{Q_i^\sharp(u)(x, s) + L_i^\sharp(u)(x, s)\} ds.$$

**Remark :** The solutions defined as above are weaker than those in the distribution sense.

**Main Theorem.**— Suppose the conditions (vd) and ( $\mu rQ$ ). Let the Cauchy data be positive with locally finite entropy (not necessary bounded). Then there exists a global solution in time and the solution is unique and positive.

Let's show the local existence in time for the small mass :

**Theorem 2.**— Suppose the condition (vd). There exists  $\delta > 0$  such that, for  $\|u^0\|_{L^1} \leq \delta$ , there exists an unique solution  $u$  in  $\mathcal{B}(\mathbf{R} \times [0, \delta])$ , and we have  $\|u\|_{\mathcal{B}} \leq 2\delta$ . Furthermore, the mapping  $u^0 \mapsto u$  is continuous from the ball with radius  $\delta$  of  $L^1$  to the ball with radius  $2\delta$  of  $\mathcal{B}(\mathbf{R} \times [0, \delta])$ . Finally, we have the finite velocity propagation and the conservation of the positivity.

**Proof.** The equation can be written in the form  $u - Ku = f$ , where

$$(Ku)_i(x, t) = \int_0^t (Q_i(u) + L_i(u))(x - c_i(t - s), s) ds$$

and  $(f)_i = u_i^0(x - c_i t)$ . For sufficiently small  $\delta$ ,  $K$  is Lipschitz continuous with Lipschitz constant  $1/2$  from the ball with radius  $\delta$  of  $L^1$  to the ball with radius  $2\delta$  of  $\mathcal{B}(\mathbf{R} \times [0, \delta])$ . We have also  $K(0) = 0$  and  $\|v\|_{\mathcal{B}} \leq \delta' \Rightarrow \|Kv\|_{\mathcal{B}} \leq \delta'/2$  for  $\delta' \leq \delta$ . Then the inverse of  $1 - K$  exists in the form of the Neumann series  $\sum_{n=0}^{\infty} K^n$  and it is Lipschitz continuous with Lipschitz constant 2. The mapping  $u^0 \mapsto u$  is continuous from the ball with radius  $\delta$  of  $L^1$  to the ball with radius  $2\delta$  of  $\mathcal{B}(\mathbf{R} \times [0, \delta])$ .

Let's show the finite velocity propagation. Now we suppose that the Cauchy data are supported in  $[-B, B]$  with  $B < \infty$ . Then it is easy to see that  $Ku$  is supported in

$\{(x, t) : x \in [-B - Ct, B + Ct]\}$  where  $C = \max_i |c_i|$ . Similarly,  $K^2 u$ , then all the  $K^n u$  ( $n \geq 1$ ), are supported in the same region. Because we have  $u = \sum_{n=0}^{\infty} K^n f$ , the support of  $u$  is included in  $\{(x, t) : x \in [-B - Ct, B + Ct]\}$ . Then we have the finite velocity propagation.

Now we suppose that  $u^0$  is bounded. Let  $[0, T]$  with  $T \leq \delta$  be the widest interval where  $\sup_{i,x} |u_i(x, \cdot)|$  is bounded. Then  $u$  is a solution in the distribution sense. It is easy to see  $T > 0$ . We put  $M = \sup_{t \in [0, T]} \sup_{i,x} |u_i(x, t)| < \infty$ . By the equation, we have

$$\begin{aligned} \sup_x u_i^\dagger(x, t) &\leq \|u^0\|_{L^\infty} + C_* M \|u\|_B + C_* \|u\|_B + MT \\ &\leq C_* \delta M + C_* \delta + \|u^0\|_{L^\infty} \end{aligned}$$

where the constant  $C_*$  depends only on the equation. For  $\delta < (2C_*)^{-1}$ , we have

$$M = \sup_{t \in [0, T]} \sup_{i,x} |u_i(x, t)| \leq 2(1 + \|u^0\|_{L^\infty}).$$

This upperbound depends only on the Cauchy data. Therefore we have  $T = \delta$ . We omit a proof for the conservation of the positivity. (See [14]) ■

**Corollary 3.**— Suppose the condition (vd). Let  $u^0$  be positive data in  $L^1$  and  $h$  a number such that  $\int_a^{a+h} u_i^0 \leq \delta$  for any  $a \in \mathbb{R}$ , then there exists an unique solution  $u$  in  $\mathcal{B}(\mathbb{R} \times [0, \theta])$  with  $\theta = \min\{\delta, h/C\}$  and  $C = \max_i |c_i|$ . Furthermore we have the finite velocity propagation. Finally, if we suppose the condition  $(\mu r Q)$ , then, for the Cauchy data such that  $u^0$  are supported in  $[-R, R]$  and verifying  $\sum_i \int_{\mathbb{R}} u_i^0 \log u_i^0(x) dx < \infty$ , we have, for  $t \in [0, \theta]$ ,  $H(t) + K \leq e^{C_* t} (H(0) + K)$  with  $C_*$  which depends only on the equation, where  $H(t) = \sum_i \int_{\mathbb{R}} u_i \log u_i(x, t) dx$  and  $K$  depends only on the equation and  $R$ .

**Proof.** By virtue of the finite velocity propagation shown in the preceding theorem, we can restrict the data in the interval  $[a, a + h]$ . Then the solution exists in small triangles of base  $[a, a + h]$  and of height  $\min\{\delta, h/C\}$ . Pasting the solutions defined in these triangles, we have a solution in  $\mathcal{B}$  up to the time  $\theta$  with  $\theta = \min\{\delta, h/C\}$ .

For the calculus on the increase of  $H(t)$ , we make approach the data by  $u_n^0 = \inf(u^0, n)$ . Then we have  $\sum_i \int_{\mathbb{R}} u_{n,i}^0 \log u_{n,i}^0(x, t) dx \leq \sum_i \int_{\mathbb{R}} u_i^0 \log u_i^0(x, t) dx < \infty$ . By virtue of the theorem 2, the solutions  $u_n$  corresponding to the Cauchy data  $u_n^0$  exist up to the time  $\theta$ . Furthermore they are bounded and positive up to the time  $\theta$ . Especially the solutions have their support in  $x$  included in  $[-R', R']$  with  $R' = R + C_* \theta$ . Therefore the quantity

$H_n(t) = \sum_i \int_{\mathbf{R}} u_{n,i} \log u_{n,i}(x, t) dx$  are well-defined. We write

$$\begin{aligned}
& \sum_i \left( \frac{\partial}{\partial t} + c_i \frac{\partial}{\partial x} \right) (u_{n,i} \log u_{n,i} e^{-Ct}) \\
&= -C u_{n,i} \log u_{n,i} e^{-Ct} \\
&- \frac{1}{2} e^{-Ct} \sum_{ijkl} A_{ij}^{kl} (u_{n,k} u_{n,\ell} \log \frac{u_{n,k} u_{n,\ell}}{u_{n,i} u_{n,j}} - u_{n,k} u_{n,\ell} + u_{n,i} u_{n,j}) \\
&- e^{-Ct} \sum_{ik} \alpha_i^k u_{n,k} (\log u_{n,k} - \log u_{n,i}) \\
&\leq -C e^{-Ct} \sum_i u_{n,i} \log u_{n,i} - e^{-Ct} \sum_{ik} \alpha_i^k u_{n,k} (\log u_{n,k} - \log u_{n,i}),
\end{aligned}$$

where  $C$  is a constant which we choose later. Integrating this inequality over  $[0, t]$  then over  $\mathbf{R}$ , we obtain

$$\begin{aligned}
& e^{-Ct} H_n(t) - H_n(0) \\
&\leq -C e^{-Ct} \int_0^t H_n(\tau) d\tau - e^{-Ct} \sum_{ik} \alpha_i^k \int_0^t \int_{\mathbf{R}} u_{n,k} \log u_{n,k} d\tau dx \\
&+ e^{-Ct} \sum_{ik} \alpha_i^k \int_0^t \int_{\mathbf{R}} u_{n,k} \log u_{n,i} d\tau dx.
\end{aligned}$$

We have

$$\begin{aligned}
& -e^{-Ct} \sum_{ik} \alpha_i^k \int_0^t \int_{\mathbf{R}} u_{n,k} \log u_{n,k} d\tau dx \leq C_* t e^{-Ct} R' \\
&\leq C_* R' t
\end{aligned}$$

with a constant  $C_*$  which depends only on the equation, because we have  $-x \log x \leq 1/e$  for any  $x \in \mathbf{R}$ . Because  $u_k \log u_i \leq \max\{0, u_i \log u_i, u_k \log u_k\}$ , we obtain

$$\begin{aligned}
& \sum_{ik} \alpha_i^k \int_0^t \int_{\mathbf{R}} u_k \log u_i d\tau dx \leq C_* e^{-Ct} \int_0^t H_n(\tau) d\tau + C_* t e^{-Ct} R' \\
&\leq C_* e^{-Ct} \int_0^t H_n(\tau) d\tau + C_* R' t
\end{aligned}$$

with a constant  $C_*$  which depends only on the equation. We obtain finally

$$e^{-Ct} H_n(t) - H_n(0) \leq -(C - C_*) e^{-Ct} \int_0^t H_n(\tau) d\tau + C_* R' t.$$

Taking the constant  $C > C_*$ , and using  $-H_n(t) \leq C_* R'$ , we have

$$H_n(t) \leq e^{Ct} (H(0) + C_* R' t).$$

Putting  $K = C_* R'$ , we have

$$\begin{aligned} H_n(t) + K &\leq e^{Ct} (H(0) + K + Kt) \\ &\leq e^{Ct} (H(0) + Ke^t) \\ &\leq e^{(C+1)t} (H(0) + K) . \end{aligned}$$

By virtue of the conservation of the mass, the  $u_n(\cdot, t)$  converge to the solution  $u(\cdot, t)$  in  $L^1$  for each  $t \in [0, \theta]$ . Taking a suitable sub-sequence, the  $u_n(\cdot, t)$  converge to  $u(\cdot, t)$  almost everywhere. Because  $u_{n,i} \log u_{n,i}(\cdot, t)$  are upper-bounded by  $1/e$  and that they are supported in a fixed compact set, the Fatou's lemma give that

$$\begin{aligned} H(t) + K &= \sum_i \int_{\mathbf{R}} u_i \log u_i(x, t) dx + K \\ &\leq \liminf \sum_i \int_{\mathbf{R}} u_{n,i} \log u_{n,i}(x, t) dx + K \\ &= \liminf H_n(t) + K \leq e^{(C+1)t} (H(0) + K) . \blacksquare \end{aligned}$$

**Corollary 4.**— Suppose the conditions (vd).

1) Suppose that there exist two solutions  $u$  and  $v$  in  $\mathcal{B}(\mathbf{R} \times [0, T])$  corresponding to the summable and positive data which coincident in an interval  $[a, b]$ . Then the solutions coincident in the triangle or trapezoid  $\{(x, t) : t \in [0, T], x \in [a + Ct, b - Ct]\}$ .

2) Let the Cauchy data be supported in  $[-R, R]$ , summable and positive. Suppose that there exists a solution  $u$  in  $\mathcal{B}(\mathbf{R} \times [0, T])$ . Then the support of  $u(\cdot, t)$  is included in  $[-R - Ct, R + Ct]$ .

**Proof.** Let be  $t_0 = \inf\{t : u(\cdot, t) \neq v(\cdot, t)\}$ . We have then  $u(\cdot, t_0) = v(\cdot, t_0) \in L^1$ . Because  $u$  and  $v$  are in  $L^1$ , there exists  $p$  such that  $\int_{\{x : u(x, t_0) \geq p\}} u(x, t_0) dx < \delta/2$ . Taking  $h = \delta/(2p)$ , we have, for any  $a \in \mathbf{R}$ ,

$$\int_a^{a+h} u_i(x, t_0) dx \leq \frac{\delta}{2} + hp \leq \delta .$$

Using the preceding corollary,  $u$  and  $v$  coincident in small triangles of base  $[a, a+h] \cap \{t = t_0\}$  and of height  $\theta$ , which is a contradiction.

To prove the finite velocity propagation, we introduce

$$t_0 = \inf\{t \text{ such that the support of } u(\cdot, t) \text{ is not included in } [-R - Ct, R + Ct] \} .$$

Then, similarly, there exists  $h$  such that  $\int_a^{a+h} u_i(x, t_0) dx \leq \delta$ . By virtue of the preceding corollary, the support of  $u(\cdot, t)$  is included in  $[-R - Ct, R + Ct]$  up to the time  $t_0 + \theta$ , which is a contradiction.  $\blacksquare$

**Lemma 5.**— Suppose the conditions (vd) and ( $\mu rQ$ ). Let  $u(x, t)$  be a positive solution defined in  $\mathbf{R} \times [0, T]$  and supported in  $[-R, R]$ . We suppose that, for any  $t \in [0, T]$ ,  $\sum_i \int_{\mathbf{R}} u_i \log u_i(x, t) dx$  is less than a constant  $C$  which don't depend on  $t$ . Then, for any  $\delta > 0$ , there exists a  $h$  which depends only on  $R, C$ , and  $\delta$  such that  $\int_a^{a+h} u(x, t) dx \leq \delta$  for any  $a \in \mathbf{R}$  and any  $t \in [0, T]$ .

**Proof.** If not, for any  $h > 0$ , there should exist  $a_* \in \mathbf{R}$  and  $t_* \in [0, T^*]$  such that  $\int_{a_*}^{a_*+h} u_i(x, t_*) dx > \delta$ . Now we use the argument due to Toscani [8] and Tartar-Crandall [7]. We put, for  $m \geq 1$ ,

$$B_1 = \{x \in [a_*, a_* + h] : u_i(x, t_*) \geq e^m\}$$

and

$$B_2 = [a_*, a_* + h] \setminus B_1.$$

Then we should have

$$\int_{a_*}^{a_*+h} u_i(x, t_*) dx \leq \frac{1}{m} \int_{B_1} u_i \log^+ u_i(y, t_*) dy + h e^m$$

where  $\log^+ y = \max\{\log y, 0\}$ . On the other hand, we should have

$$\begin{aligned} C &\geq \sum_i \int_{\mathbf{R}} u_i \log u_i(y, t_*) dy \\ &\geq \sum_i \int_{-R}^R (u_i \log^+ u_i(y, t_*) - 1) dy \\ &= \sum_i \int_{-R}^R u_i \log^+ u_i(y, t_*) dy - 2pR, \end{aligned}$$

where  $p = \#I$ . Then we should obtain

$$\begin{aligned} \delta &< \sum_i \int_{a_*}^{a_*+h} u_i(y, t_*) dy \\ &\leq \frac{1}{m} \sum_i \int_{a_*}^{a_*+h} u_i \log^+ u_i(y, t_*) dy + p h e^m \\ &\leq \frac{1}{m} (C + 2pR) + p h e^m. \end{aligned}$$

Choosing  $m$  such that  $\frac{1}{m}(C + 2pR) < \frac{\delta}{4}$ , then  $h$  such that  $p h e^m < \frac{\delta}{4}$ , we should have  $\delta < \frac{\delta}{4} + \frac{\delta}{4} = \frac{\delta}{2}$ , which is a contradiction. ■

**Corollary 6.**— Suppose the conditions (vd) and ( $\mu$ rQ). Let the Cauchy data be supported in  $[-R, R]$ , positive with finite entropy. Suppose that there exists a solution in  $\mathcal{B}(\mathbf{R} \times [0, T])$ . Then we have

$$H(t) + K \leq e^{C_* t} (H(0) + K)$$

where the constant  $C_*$  depends only on the equation and  $K$  depends only on the equation,  $R$  and  $T$ .

**Proof.** The solution have its support in  $x$  included in  $[-R', R']$  with  $R' = R + CT$ . Let  $K$  be the constant associated to  $R'$  by the corollary 3. Let be  $t_0 = \inf\{t \text{ such that the estimate is not valid at the time } t\}$ . Taking a small  $\varepsilon > 0$ , we have  $H(t_0 - \varepsilon) \leq e^{C_*(t_0 - \varepsilon)} (H(0) + K) \leq e^{C_* T} (H(0) + K)$ . Because  $u(\cdot, t_0 - \varepsilon)$  is positive and supported in  $[-R', R']$ , the preceding lemma shows that there exists a  $h$  independent of  $\varepsilon$  such that we have, for any  $a \in \mathbf{R}$ ,  $\int_a^{a+h} u_i(x, t_0 - \varepsilon) dx \leq \delta$ . By virtue of the corollary 3, we have

$$\begin{aligned} H(t_0 - \varepsilon + \theta) + K &\leq e^{C_* \theta} (H(t_0 - \varepsilon) + K) \\ &\leq e^{C_* \theta} e^{C_*(t_0 - \varepsilon)} (H(0) + K). \end{aligned}$$

The estimate is then verified up to the time  $t_0 - \varepsilon + \theta$  with  $\theta > 0$  independent of  $\varepsilon$ , which is a contradiction. ■

Now let's show the proof of the main theorem.

**Proof.** Let  $T_0$  be arbitrary number and we want to prove that the existence time of the solution is at least  $T_0$ . By virtue of the finite velocity propagation, we can suppose that the Cauchy data are supported in  $[-R', R']$  and they have a finite entropy. Let  $T^*$  be the existence time of the solution. Suppose that  $T^* < T_0$ . By the corollary 6, the entropy is bounded for  $t < T^*$ :  $H(t) \leq H_T < \infty$ . By virtue of the lemma 5, there exists  $h > 0$  such that  $\int_a^{a+h} u_i(x, t) dx \leq \delta$  for any  $a \in \mathbf{R}$  and  $t < T^*$ . For  $t < T^*$ , by the corollary 3, applied to the data  $u_i(\cdot, t)$ , there exists  $\theta > 0$  independent of  $t$  such that the solution is prolonged in  $\mathbf{R} \times [0, t + \theta]$ . It is sufficient to choose  $t > T^* - \theta$  for arriving at a contradiction. ■

## REFERENCES

1. Bony J.M., *Solutions globales bornées pour les modèles discrets de l'équation de Boltzmann, en dimension 1 d'espace*, Actes des journées E.D.P. à Saint-Jean-de-Monts (1987), Ecole Polytechnique.
2. Bony J.M., *Problème de Cauchy et diffusion à données petites pour les modèles discrets de la cinétique des gaz*, Actes des journées E.D.P. à Saint-Jean-de-Monts (1990), Ecole Polytechnique.
3. Bony J.M., *Existence globale et diffusion en théorie cinétique discrète*, Actes du Symposium en l'honneur de Henri Cabannes, Paris (1990).
4. Bony J.M., *Existence globale à données de Cauchy petites pour les modèles discrets de l'équation de Boltzmann*, Comm. in Partial Differential Equations 16 (1991), 533-545.
5. Carleman T., *Problèmes mathématiques dans la théorie cinétique des gaz*, Uppsala (1957), Scientific Publications of Mittag-Leffler Institute.
6. Nishida T., Mimura M., *On the Broadwell's model for a simple discrete velocity gas*, Proc. Japan Acad. 50 (1974), 812-817.
7. Tartar L., Crandall M.G., *Existence globale pour un système hyperbolique de la théorie des cinétique des gaz*, Ecole Polytechnique, Séminaire Goulaouic-Schwartz n°1 (1975).



8. Toscani G., *On the Cauchy problem for the discrete Boltzmann equation with initial value in  $L^1_+(\mathbf{R})$* , Comm.Math.Phys. (1989), 121–142.
9. Yamazaki M., *Existence globale pour les modèles discrets de l'équation de Boltzmann dans un tube mince infini*, C.R.Acad.Sci., Série I, Paris 313 (1991), 29–32.
10. Yamazaki M., *Existence globale pour les modèles discrets de l'équation de Boltzmann dans un tube mince infini en dimension 1 d'espace*, J.Fac.Sci.Univ.Tokyo, Sec. IA 39 (1992), 43–59.
11. Yamazaki M., *Etude de l'équation de Boltzmann pour les modèles discrets dans un tube mince infini*, Transport Theory and Statistical Physics, Proceedings of the Fourth MAFPD 21 (1992), 465–473.
12. Yamazaki M., *On the Discrete Boltzmann Equation with Linear and Nonlinear Terms*, Thesis at University of Tokyo.
13. Yamazaki M., *On the Discrete Boltzmann Equation with Linear and Nonlinear Terms*, J.Fac.Sci.Univ. Tokyo (to appear).
14. Yamazaki M., *Sur les modèles discrets de l'équation de Boltzmann avec termes linéaires et quadratiques*, Thesis of Ecole Polytechnique, Palaiseau, France.